

Existence and Uniqueness Solution of Fractional Integral Equations of Volterra Types

Sebar T. Qader ¹, Samir H. Abbas ²

^{1,2}Department of Mathematics, Collage of Basic Education, University of Duhok, Kurdistan Region, Iraq

ABSTRACT

One of the objectives of this paper is to study the existence, uniqueness and stability solutions of Fractional integral equations by using both methods Picard approximation and Banach fixed point theorems. This paper guides us to develop the Picard approximation and Banach fixed point theorems and to expand the results got by Butris.

Keywords: Picard approximation method, Banach fixed point theorem, existence, uniqueness and stability solutions.

1. Introduction

1.1. Brief history review

Many results about the existence, uniqueness and stability solutions of Fractional integral equations have been obtain by Picard approximation method and Banach fixed point theorem that were proposed by [7] which had been later applied in many studies [2,4,5,6,8]. Fractional calculus is a branch of mathematical analysis that studies the several different possibilities of defining real number powers or complex number powers of the differentiation operator and integration operator.

Riemann in 1847 while a student wrote a paper published posthumously in which he gives a definition of a fractional operation. Riemann was influenced by

one of Liouville's memoirs in which Liouville wrote, "The ordinary differential equation has the complementary solution. Thus should have a corresponding complementary solution." Then Riemann saw fit to add a complementary function to his definition of a fractional integration [3]. Butris [1] used both methods Picard approximation and Banach fixed point theorems for studying the existence and uniqueness solutions to the following fractional integral equations: -

$$\begin{aligned} u(t) &= u_0 + f(t) \\ &+ \frac{1}{\Gamma(\alpha)} \int_t^{\infty} (t-s)^{\alpha-1} k(t,s) F(s, u(s), w(s)) ds, \quad (t \in [a, b]) \end{aligned}$$

$$\begin{aligned} w(t) &= w_0 + g(t) \\ &+ \frac{1}{\Gamma(\alpha)} \int_t^{\infty} (t-s)^{\alpha-1} H(t,s) G(s, u(s), w(s)) ds \end{aligned}$$

Academic Journal of Nawroz University (AJNU)
Volume 8, No 4 (2019).

Regular research paper : Published 28 Dec 2019

Corresponding author's e-mail :
raad.butris@gmail.com

Copyright ©2018 iRaad N. Butris, 2Sebar T. Qader.
This is an open access article distributed under the
Creative Commons Attribution License.

In this equation the functions $f(t)$ and $F(t, s)$ be continuous on (finite) interval $a \leq t \leq b$ and the square region $a \leq t \leq b, a \leq s \leq b$, respectively.

Our work is extend some results of Butris [1] to Volterra types, by using also both methods.

Consider the following integrals equations of Volterra types

$$\begin{aligned} u(t) &= u_0 + f(t) \\ &+ \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t (t-s)^{\alpha-1} K(t, s) F(s, u(s), w(s)) ds \dots \dots \dots (V_1) \end{aligned}$$

and

$$\begin{aligned} w(t) &= w_0 + g(t) \\ &+ \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t (t-s)^{\alpha-1} H(t, s) G(s, u(s), w(s)) ds \dots \dots (V_2) \end{aligned}$$

where

$u \in D_1 \subseteq \mathbb{R}^n, w \in D_2 \subseteq \mathbb{R}^m, D_1, D_2$ are a compact domains.

The vector functions $f(t), g(t)$ and $F(t, u, w), G(t, u, w)$ are define and continuous on the domain

$$\begin{aligned} D &= \{(t, u, w); t \in \mathbb{R}^1, u \in D_1, w \in D_2\}. \dots \dots \dots (1) \end{aligned}$$

Suppose that the vector functions $F(t, u, w)$ and $f(t, u, w)$ satisfy the following inequalities

$$\left. \begin{aligned} \|F(t, u, w)\| &\leq M_1, \\ \|G(t, u, w)\| &\leq M_2 \\ \|F(t, u_1, w_1) - F(t, u_2, w_2)\| &\leq K_1 \|u_1 - u_2\|^\gamma + K_2 \|w_1 - w_2\|^\gamma \\ \|G(t, u_1, w_1) - G(t, u_2, w_2)\| &\leq L_1 \|u_1 - u_2\|^\beta + L_2 \|w_1 - w_2\|^\beta \end{aligned} \right\} \dots (2)$$

For all $t \in \mathbb{R}^1, u, u_1, u_2 \in D_1, w, w_1, w_2 \in D_2$ where M_1, M_2, K_1, K_2 and L_1, L_2 are

positive constant provided that

$$\|K(t, s)\| \leq \delta_1 e^{-\lambda_1(t-s)}, \|H(t, s)\| \leq \delta_2 e^{-\lambda_2(t-s)}, \|t - s\|^{\alpha-1} \leq b^{\alpha-1} \dots (3)$$

where $\delta_1, \delta_2, \lambda_1, \lambda_2 > 0$, with $\|\cdot\| = \max_{t \in [a, b]} |\cdot|$.

We define non-empty sets as follows: -

$$\left. \begin{aligned} D_F &= D_1 - \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_1}{\lambda_1} M_1 \|u_{m-1} - u_0\|^\gamma \\ D_G &= D_2 - \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_2}{\lambda_2} M_2 \|u_{m-1} - u_0\|^\beta \end{aligned} \right\} \dots (4)$$

Furthermore, we assume that the largest Eigen-value of the matrix

$$Q = \begin{pmatrix} K_1 \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_1}{\lambda_1} b \|u_{m-1} - u_0\|^{\gamma-\lambda} \\ K_2 \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_1}{\lambda_1} b \|u_{m-1} - u_0\|^\gamma \\ L_1 \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_2}{\lambda_2} b \|u_{m-1} - u_0\|^\beta \\ L_2 \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_2}{\lambda_2} b \|u_{m-1} - u_0\|^{\beta-\lambda} \end{pmatrix}$$

greater than zero and less than one

$$\text{i.e } 0 < \lambda \max(Q) < 1 \dots \dots \dots (5)$$

$$\text{where } \lambda_{\max} = \frac{(\mu+\zeta) \pm \sqrt{(\mu+\zeta)^2 - 4(\mu\zeta - \xi\eta)}}{2}$$

$$\mu = K_1 \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} \|u_{m-1} - u_0\|^\gamma, \quad \xi =$$

$$\eta = L_1 \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} \|u_{m-1} - u_0\|^\beta,$$

$$\zeta = L_2 \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} \|u_{m-1} - u_0\|^\beta.$$

Define a sequence of functions $\{u_m(t), w_m(t)\}_{m=0}^\infty$ by the following

$$u_m(t) = u_0 + f(t) + \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t (t-s)^{\alpha-1} K(t,s) F(s, u_{m-1}(s), w_{m-1}(s)) ds$$

with

$$u_0(0) = u_0 + f(t) \dots \dots \dots (6)$$

and

$$w_m(t) = w_0 + g(t) + \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t (t-s)^{\alpha-1} H(t,s) H(s, u_{m-1}(s), w_{m-1}(s)) ds$$

with

$$w_0(0) = w_0 + g(t) \dots \dots \dots (7)$$

1.2. Preliminaries

In this section, we give some definitions and theorems which are needed in this work.

Definition 1.[1].

Let $(D, \|\cdot\|)$ be normed space if T maps D into itself, we say that T is a contraction mapping on D if there exists $\alpha \in \mathbb{R}^1$ with $0 < \alpha < 1$ such that $\|Tx(t) - Ty(t)\| \leq \alpha \|x(t) - y(t)\|, (x(t), y(t)) \in D$.

$$K_2 \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} \|u_{m-1} - u_0\|^\gamma,$$

Definition 2.[1] (Lipchitz Condition)

Let f be a continuous function defined on a domain $D = \{(t, x): a \leq t \leq b, c \leq x \leq d\}$. Then f is said to satisfy a Lipchitz condition in the variable x on D , provided that a constant $L > 0$ exists with property that $|f(t, x_1) - f(t, x_2)| \leq L|x_1 - x_2|$, for all $(t, x_1), (t, x_2) \in D$. the constant L is called a Lipchitz constant for f .

Definition 3. [8] (Stability)

A solution $x(t)$ is said to be stable if for each $\varepsilon > 0$, there exists a $\delta > 0$ such that any solution $\check{x}(t)$ which satisfies $\|\check{x}(t_0) - x(t_0)\| < \delta$ for some t_0 , also satisfies $\|\check{x}(t) - x(t)\| < \varepsilon$ for all $t \geq t_0$.

Definition 4. [8]

Let $\dot{x} = f(t, x)$ be a system of differential equation and let $x(t)$ be a solution defined for $t \geq t_0$. The solution $x(t)$ is said to be stable if for each $\varepsilon > 0$ there exists a positive $\delta = \delta(\varepsilon)$, such that any solution $\bar{x}(t)$ which is defined at $t = t_0$ and satisfied $\|\bar{x}(t_0) - x(t_0)\| \leq \delta$ is defined for all $t \geq t_0$ and satisfies $\|\bar{x}(t) - x(t)\| < \varepsilon$ for all $t \geq t_0$.

Definition 5. [1]

Let $\{f_m(t)\}_{m=0}^\infty$ be a sequence of functions f defined on a set $E \subseteq \mathbb{R}^1$ we say that $\{f_m(t)\}_{m=0}^\infty$ converges uniformly to the limit function f on E if given $\varepsilon > 0$ there exists a positive integer such that $|f_m(t) - f(t)| < \varepsilon$,

$(m \geq N, t \in E)$.

Theorem 1. [1] (Banach fixed point theorem)

E be a Banach space. If T is a contraction mapping on E, then T has one and only fixed point in E.

Theorem 2. [1]

Let $c[0, h]$ be a space of all continuous functions on $[0, h]$, for all $x(t) \in c[0, h]$, we denoted $\|x(t)\| = \max_{t \in [0, h]} |x(t)|$.

Then $(c[0, h], \|\cdot\|)$ is a Banach space.

1. Existence Solutions of (V_1) and (V_2) .

The investigation of the existence solution of (V_1) and (V_2) will be the introduced by the following theorem:-

Theorem4. Let the vector functions $f(t), g(t)$ and $F(t, u, w), G(t, u, w)$ defined and continuous on the domain (1) suppose these functions are satisfying the inequalities (2), (3) and the conditions (4), (5). Then there exist a sequence of functions (6) and (7) converges uniformly on the domain

$$\begin{aligned} G^* &= \{(t, u_0, w_0) \\ &\in [a, b] \times D_1 \\ &\times D_2\} \dots \dots \dots (8) \end{aligned}$$

to the limit vector function $\begin{pmatrix} u(t) \\ w(t) \end{pmatrix}$ which is a continuous on the domain (1) and satisfies the following integral equations:-

$$\begin{pmatrix} u(t) \\ w(t) \end{pmatrix} = \begin{pmatrix} u_0 + f(t) + \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t (t-s)^{\alpha-1} K(t, s) F(s, u_0(s), w_0(s)) ds \\ w_0 + g(t) + \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t (t-s)^{\alpha-1} H(t, s) G(s, u_0(s), w_0(s)) ds \end{pmatrix} \dots \dots \dots (9)$$

and it is exist solution of (V_1) and (V_2) .

Provide that

$$\begin{pmatrix} \|u_m(t) - u_0\| \\ \|w_m(t) - w_0\| \end{pmatrix} \leq \begin{pmatrix} \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_1}{\lambda_1} M_1 \|u_{m-1} - u_0\|^\gamma \\ \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_2}{\lambda_2} M_2 \|u_{m-1} - u_0\|^\beta \end{pmatrix} \dots (10)$$

and

$$\begin{pmatrix} \|u_{m+1}(t) - u_0(t)\| \\ \|w_{m+1}(t) - w_0(t)\| \end{pmatrix} \leq Q^m (E - Q)^{-1} \Psi_0 \dots \dots \dots (11)$$

for all $t \in [a, b]$ and $u_0 \in D_F, w_0 \in D_G, m = 0, 1, 2, \dots$

where

$$\Psi_0 = \begin{pmatrix} \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_1}{\lambda_1} M_1 \|u_{m-1} - u_0\|^\gamma \\ \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_2}{\lambda_2} M_2 \|u_{m-1} - u_0\|^\beta \end{pmatrix}$$

By mathematical indication we can prove that

$$\begin{aligned} &\|u_m(t) - u_0\| \\ &\leq \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_1}{\lambda_1} M_1 \|u_{m-1} - u_0\|^\gamma \\ &\dots \dots \dots (12) \end{aligned}$$

That is $u_m(t) \in D_F$, for all $t \in [a, b]$ and $u_0 \in D_F$.

Similarly, from the sequence of functions (7) and we obtain that

$$\begin{aligned} & \|w_m(t) - w_0\| \\ & \leq \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} M_2 \|u_{m-1} - u_0\|^\beta \\ & \quad \dots \dots \dots (13) \end{aligned}$$

That is $w_m(t) \in D_G$, for all $t \in [a, b]$ and $w_0 \in D_G$.

Next we shall prove that the sequences of functions (6) and (7) converge uniformly on the domain (8). Then by mathematics induction we have

$$\begin{aligned} \|u_{m+1}(t) - u_m(t)\| & \leq \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} M_1 \|u_{m-1} - u_0\|^\gamma (K_1 \|u_m(t) - u_{m-1}(t)\|^\gamma + \\ & K_2 \|w_m(t) - w_{m-1}(t)\|^\gamma) \dots \dots \dots (14) \end{aligned}$$

and

$$\begin{aligned} \|w_{m+1}(t) - w_m(t)\| & \leq \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} M_2 \|u_{m-1} - u_0\|^\beta (L_1 \|u_m(t) - u_{m-1}(t)\|^\beta + \\ & L_2 \|w_m(t) - w_{m-1}(t)\|^\beta) \dots \dots \dots (15) \end{aligned}$$

Rewrite (14) and (15) in a vector form we get

$$\begin{aligned} \Psi_{m+1} & \leq Q(t) \\ & \leq \Psi_m \dots \dots \dots (16) \end{aligned}$$

where

$$\Psi_{m+1} = \begin{pmatrix} \|u_{m+1}(t) - u_m(t)\| \\ \|w_{m+1}(t) - w_m(t)\| \end{pmatrix}$$

$$\Psi_m = \begin{pmatrix} \|u_m(t) - u_{m-1}(t)\| \\ \|w_m(t) - w_{m-1}(t)\| \end{pmatrix},$$

and

$$Q(t) = \begin{pmatrix} K_1 \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} t \|u_{m-1} - u_0\|^\gamma \\ K_2 \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} t \|u_{m-1} - u_0\|^\gamma \\ L_1 \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} t \|u_{m-1} - u_0\|^\beta \\ L_2 \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} t \|u_{m-1} - u_0\|^\beta \end{pmatrix}$$

Now we take the maximum value for the both sides of the inequality (16)

$$\begin{aligned} \Psi_{m+1} & \leq Q \Psi_m \dots \dots \dots (17) \end{aligned}$$

where $Q = \max_{t \in [a, b]} Q(t)$.

By repetition of (17) we find that $\Psi_{m+1} \leq Q^m \Psi_1$ and also we get

$$\sum_{i=1}^m \Psi_i \leq \sum_{i=1}^{\infty} Q^{i-1} \Psi_0 \dots \dots \dots (18)$$

Using the condition (15), thus the sequence of functions (6) and (7) are uniformly convergent, that is

$$\begin{aligned} & \lim_{m \rightarrow \infty} \sum_{i=1}^m Q^{m-1} \Psi_0 \\ & = \sum_{i=1}^{\infty} Q^{m-1} \Psi_0 \\ & = (1 - Q)^{-1} \Psi_0 \dots \dots \dots (19) \end{aligned}$$

Let

$$\begin{aligned} & \lim_{m \rightarrow \infty} \begin{pmatrix} u_m(t) \\ w_m(t) \end{pmatrix} \\ & = \begin{pmatrix} u(t) \\ w(t) \end{pmatrix} \dots \dots \dots (20) \end{aligned}$$

Since the sequence of function (6) and (7) are defined and continuous in the domain (1) then the limit vector function

$\begin{pmatrix} u(t) \\ w(t) \end{pmatrix}$ is also defined and continuous in the domain (1).

By using the conditions and inequalities of a theorem, we can prove that the inequalities (10) and (11) will be satisfied for all $t \in [a, b]$, $u_0 \in D_F$, $w_0 \in D_G$, $m = 0, 1, 2, \dots$.

2. Uniqueness Solutions of (V_1) and (V_2) .

The investigation of the uniqueness solutions of (V_1) and (V_2) will be introduced by :-

Theorem 5. Let all assumptions and conditions of Theorem 3 be satisfied.

Then the solution $\begin{pmatrix} u(t) \\ w(t) \end{pmatrix}$ is a unique of (V_1) and (V_2) .

Proof. Let $\begin{pmatrix} \check{u}(t) \\ \check{w}(t) \end{pmatrix}$ be another solution of (V_1) and (V_2) , that is

$$\begin{aligned} \check{u}(t) = & u_0 + f(t) \\ & + \frac{1}{\Gamma(\alpha)} \int_t^t (t-s)^{\alpha-1} K(t, s) \\ & + F(s, \check{u}(s), \check{w}(s)) ds \end{aligned}$$

and

$$\begin{aligned} \check{w}(t) = & w_0 + g(t) \\ & + \frac{1}{\Gamma(\alpha)} \int_t^t (t-s)^{\alpha-1} H(t, s) \\ & + G(s, \check{u}(s), \check{w}(s)) ds \end{aligned}$$

Assuming

$$\|u(t) - \check{u}(t)\| \leq \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_1}{\lambda_1} t (K_1 \|u(s) - \check{u}(s)\|^\gamma + K_2 \|w(s) - \check{w}(s)\|^\gamma) \dots \dots (21)$$

and

$$\|w(t) - \check{w}(t)\| \leq \frac{b^{\alpha-1}}{\Gamma(\alpha)} \frac{\delta_2}{\lambda_2} t (L_1 \|u(t) - \check{u}(t)\|^\beta + L_2 \|w_m(t) - \check{w}(t)\|^\beta) \dots \dots (22)$$

$$\begin{aligned} & \left(\begin{array}{c} \|u(t) - \check{u}(t)\| \\ \|w(t) - \check{w}(t)\| \end{array} \right) \leq \\ & Q \left(\begin{array}{c} \|u(t) - \check{u}(t)\|^\gamma \\ \|w(t) - \check{w}(t)\|^\beta \end{array} \right) \dots \dots \dots (23) \end{aligned}$$

By iterating the inequality (21) we have

$$\begin{aligned} & \left(\begin{array}{c} \|u(t) - \check{u}(t)\| \\ \|w(t) - \check{w}(t)\| \end{array} \right) \\ & \leq Q^m \left(\begin{array}{c} \|u(t) - \check{u}(t)\|^\gamma \\ \|w(t) - \check{w}(t)\|^\beta \end{array} \right) \end{aligned}$$

Then by the condition (15), we find that

$$\left(\begin{array}{c} \|u(t) - \check{u}(t)\| \\ \|w(t) - \check{w}(t)\| \end{array} \right) \rightarrow \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\text{Thus } \begin{pmatrix} u(t) \\ w(t) \end{pmatrix} = \begin{pmatrix} \check{u}(t) \\ \check{w}(t) \end{pmatrix}.$$

Hence the solutions $\begin{pmatrix} u(t) \\ w(t) \end{pmatrix}$ of (V_1) and (V_2) is a unique on the domain (1).

V. Stability Solutions of (V_1) and (V_2) .

In this section, we can study the stability solutions of Volterra integral equations

(V_1) and (V_2) respectively.

Theorem 6. Suppose that the functions $F(t, u, w)$ and $f(t, u, w)$ be continuous in the domain (1) and satisfy the inequalities (2) and (3). Then the solution (9) is stable for all $t \geq 0$.

Proof: Taking

$$\|u(t) - \bar{u}(t)\| \leq \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} t \|u_0 - \bar{u}_0\|^\gamma + \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} M_1 [K_1 \|u(t) - \bar{u}(t)\|^\gamma + K_2 \|w(t) - \bar{w}(t)\|^\gamma] \dots (24)$$

and

$$\|w(t) - \bar{w}(t)\| \leq \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} t \|w_0 - \bar{w}_0\|^\beta + \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} M_2 [L_1 \|u(t) - \bar{u}(t)\|^\beta + L_2 \|w(t) - \bar{w}(t)\|^\beta] \dots (25)$$

where

$$\bar{u} = u_0 + f(t) + \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t K(t, s) + F(s, \bar{u}(s), \bar{w}(s)) ds$$

and

$$\bar{w} = w_0 + g(t) + \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t H(t, s) + G(s, \bar{u}(s), \bar{w}(s)) ds$$

Rewrite (24) and (25) in a vector form that is

$$\begin{pmatrix} \|u(t) - \bar{u}(t)\| \\ \|w(t) - \bar{w}(t)\| \end{pmatrix} \leq \begin{pmatrix} \|u_0 - \bar{u}_0\|^\gamma \\ \|w_0 - \bar{w}_0\|^\beta \end{pmatrix} + Q \begin{pmatrix} \|u(t) - \bar{u}(t)\|^\gamma \\ \|w(t) - \bar{w}(t)\|^\beta \end{pmatrix}$$

For $\|u_0 - \bar{u}_0\| \leq \delta_1, \|w_0 - \bar{w}_0\| \leq \delta_2$ then

$$\begin{pmatrix} \|u(t) - \bar{u}(t)\| \\ \|w(t) - \bar{w}(t)\| \end{pmatrix} \leq \begin{pmatrix} \delta_1 \\ \delta_2 \end{pmatrix} + Q \begin{pmatrix} \|u(t) - \bar{u}(t)\|^\gamma \\ \|w(t) - \bar{w}(t)\|^\beta \end{pmatrix}$$

By using the condition () we have

$$\begin{pmatrix} \|u(t) - \bar{u}(t)\| \\ \|w(t) - \bar{w}(t)\| \end{pmatrix} \leq \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \end{pmatrix}, \quad \epsilon_1, \epsilon_2 \geq 0.$$

Also by using the definition of the stability [] we find that $\begin{pmatrix} \bar{u}(t) \\ \bar{w}(t) \end{pmatrix}$ is a stable solution $t \geq 0$ of (V_1) and (V_2) .

3. Another results of the solutions (V_1) and (V_2) .

In this section, we can study the solution of Volterra integrals equations (V_1) and (V_2) by using:-

Theorem7. With the hypotheses and all conditions of theorem (1) the solutions of integral equations (V_1) and (V_2) are a unique on the domain (1).

Proof: Assume that $(C[a, b], \|\cdot\|)$ is a Banach spaces. Define a mapping T on $C[a, b]$ by :-

$$\begin{aligned} Tu^*(t) &= u_0 + f(t) \\ &+ \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t K(s, t) F(s, u(s), w(s)) ds \end{aligned}$$

and

$$\begin{aligned} Tw^*(t) &= w_0 + g(t) \\ &+ \frac{1}{\Gamma(\alpha)} \int_{-\infty}^t H(s, t) G(s, u(s), w(s)) ds \end{aligned}$$

then

$$\begin{aligned} \|Tu(t) - Tu^*(t)\| &\leq \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} b (K_1 \|u(t) - u^*(t)\|^\gamma \\ &+ K_2 \|w(s) - w^*(t)\|^\gamma) \\ \dots (27) \end{aligned}$$

And

$$\begin{aligned} & \| Tw(t) - Tw^*(t) \| \\ & \leq \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} b (L_1 \| u(t) - u^*(t) \|^\beta \\ & + L_2 \| w(s) - w^*(t) \|^\beta) \\ & \dots\dots\dots(28) \end{aligned}$$

Rewrite (27) and (28) in a vector form

$$\begin{pmatrix} \| Tu(t) - Tu^*(t) \| \\ \| Tw(t) - Tw^*(t) \| \end{pmatrix} \leq \begin{pmatrix} K_1 \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} b & K_2 \frac{b^{\alpha-1} \delta_1}{\Gamma(\alpha) \lambda_1} b \\ L_1 \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} b & L_2 \frac{b^{\alpha-1} \delta_2}{\Gamma(\alpha) \lambda_2} b \end{pmatrix} \begin{pmatrix} \| u(t) - u^*(t) \|^\gamma \\ \| w(s) - w^*(t) \|^\beta \end{pmatrix}$$

By using the condition ()

Then $\begin{pmatrix} Tu^*(t) \\ Tw^*(t) \end{pmatrix}$ is a contraction mapping on $[a,b]$. By using Banach fixed point theorem such that

$$\begin{pmatrix} Tu^*(t) \\ Tw^*(t) \end{pmatrix} = \begin{pmatrix} Tu(t) \\ Tw(t) \end{pmatrix} \text{ is a fixed point on } C[a,b]$$

Then

$\begin{pmatrix} Tu^*(t) \\ Tw^*(t) \end{pmatrix}$ is exist and unique solution of (V_1) and (V_2) .

Note. Picard approximation method gives us a global solution for Volterra integral equations (V_1) and (V_2) . While contraction mapping theorem gives us a

local solutions for Voltterra integrals equations (V_1) and (V_2) .

References

- [1]. R. N. Butris , Solutions for the Volterra integral equations of the second kind, university of Mosul ,Mosul, Iraq, (1984) .
- [2]. R. N. Butris, and Rafeq, A. Sh., Existence and Uniqueness Solution for Nonlinear Volterra Integral Equation, J. Duhok Univ. Vol. 14, No. 1, (Pure and Eng. Sciences),(2011),25-29.
- [3] C. Corduneanu, I.W. Sanberg, Volterra Equations and Aplications, Gordon and Breach Science Publisher. Vol. 10, (2000) ,xiii pages.
- [4]. E. A . Coddington , An introduction to Ordinary Differential Equations ,Prentice-Hall,Inc.Englewood Cliffs,N.J., (1961), 292 pages.
- [5]F,G Tricomi, (1965), Integral Equations, Turin University, Turin, Italy.
- [6]. E.A. Goma, Sccessives approximation method of two points boundary problems, Ukrania,Math.,J.,Tom(33),Kiev,(1976),80 0-835.
- [7]. M.M. Rama, Ordinary Differential Equations Theory and Applications, United Kingdom, (1981), 266 pages.
- [8]. O.Plaat, Ordinary Differential Equations, Holden-Day,Inc.San Francisco,Cambridge,London,A msterdam ,1971)295 pages .